

Stable transformation of the moss *Physcomitrella patens*

D. Schaefer¹, J.-P. Zryd², C.D. Knight² and D.J. Cove²

¹ Laboratoire de Phytogénétique Cellulaire, Université de Lausanne, CH-1015, Switzerland

² Department of Genetics, University of Leeds, LS2 9JT, UK

Received October 31, 1990

Summary. We report the stable transformation of *Physcomitrella patens* to either G418 or hygromycin B resistance following polyethylene glycol-mediated direct DNA uptake by protoplasts. The method described in this paper was used successfully in independent experiments carried out in our two laboratories. Transformation was assessed by the following criteria: selection of antibiotic-resistant plants; mitotic and meiotic stability of phenotypes after removal of selective pressure and stable transmission of the character to the offspring. Southern hybridisation analysis of genomic DNA to show integration of the plasmid DNA; segregation of the resistance gene following crosses with antibiotic-sensitive strains; and finally Southern hybridisation analysis of both resistant and sensitive progeny. In addition to stable transformants, a heterogeneous class of unstable transformants was obtained.

Key words: *Physcomitrella patens* – Transformation

Introduction

We have chosen to use the moss *Physcomitrella patens* as a model for studying the molecular and cellular basis of development. As a model organism, *P. patens* presents the opportunity to study processes of greater morphogenetic complexity than those shown by such algae as *Chlamydomonas reinhardtii* (Snell 1985), but yet is simpler than higher plants such as *Arabidopsis thaliana* (Meyerowitz 1989). It is easily manipulated in vitro and can be taken through its life-cycle from a single spore or protoplast on a simple mineral medium within 2 to 3 months (Ashton and Cove 1977; Knight et al. 1988). Moreover, as a haploid plant, mutant isolation is facilitated and numerous biochemical and morphological mutants have already been described (Ashton and Cove 1977; Ashton et al. 1988). Genetic complementation analysis of some mutant classes using hybrids produced following protoplast fusion (Grimsley et al. 1977a) has

Offprint requests to: J.-P. Zryd

indicated the minimum number of genes involved in certain biochemical pathways or necessary for certain stages of development (Grimsley et al. 1977b; Ashton et al. 1988; Featherstone et al. 1990; C.D. Knight, submitted). In combination with the genetic data, physiological analyses of mutants and the wild-type strain have shown that the plant growth regulators, cytokinin and auxin, are required for cell differentiation both at the single cell level and for the formation of multicellular gametophores (Ashton et al. 1979a, b; Cove and Ashton 1984).

Further similarities are seen with higher plants in that most of the abiotomorphogenic processes are mediated by phytochrome (Cove et al. 1978) including protoplast regeneration (Jenkins and Cove 1983a; D. Schaefer, unpublished data), phototropism and polarotropism of protonemata, gametophores (Jenkins and Cove 1983b) and protoplasts (D. Schaefer, unpublished data).

At the molecular level, the sequence of the chlorophyll *a/b*-binding protein gene has been determined and compared with that of higher plants and the alga *Dunaliella salina* (Long et al. 1989). The *P. patens* sequence bears an intron containing enhancer-like elements and 5' sequences that are partially homologous to light-inducible sequences of higher plants. Ribulose biphosphate carboxylase, small subunit (*rbcS*) genes have also been isolated from a genomic library, using wheat and petunia cDNA probes and are being characterised (D. Schaefer, unpublished data). The restriction pattern and gene order of the chloroplast genome has also been shown to be similar to the consensus land plant genome typified by that of spinach (Culie and Hughes 1987). The continued study of morphogenesis requires analysis at the molecular level and for this purpose a transformation protocol is required. In this paper, we present the first demonstration of stable transformation of *Physcomitrella patens*.

Materials and methods

***P. patens* culture.** Culture conditions for *P. patens* are described by Ashton and Cove (1977) and Knight et al. (1988). Protoplasts were isolated according to Grimsley

et al. (1977a) by treating 5- to 7-day-old protonemata with 1%–1.4% (w/v) Driselase (Sigma Chemical Co., Dorset, UK) in 0.44 M mannitol solution for 30–60 min. In Leeds, protoplasts were regenerated according to Knight et al. (1988) in a 2.5-ml layer of medium containing 0.6% (w/v) agar (Oxoid no. 1) and 0.44 M mannitol per 9 cm petri dish, which was separated from a base layer of 1.2% (w/v) agar, 0.33 M mannitol by an 8-cm diameter cellophane disc (W.E. Canning, Bristol, UK). After 4–8 days incubation in continuous white light (15–20 W/m²) at 25°C, during which time the regenerating plants reach a stage of 5 to 10 cells, the cellophane overlays were transferred to medium without mannitol and containing the appropriate antibiotic.

In Lausanne, protoplasts were either cultivated in liquid medium for 5 days, and then embedded in a top layer of medium containing 0.6% (w/v) SeaPlaque agarose and regenerated according to Grimsley et al. (1977a) or were directly embedded in a thin layer of 0.6% (w/v) agarose in a 9 cm petri dish and regenerated in beads as described by Shillito et al. (1983). For each treatment, the appropriate antibiotic was applied after 5–7 days. The light regime in Lausanne was 16 h light, 8 h darkness. Cove et al. (1978) showed that continuous light did not adversely affect the morphological pattern of growth and we did not detect any significant difference in transformation frequencies between the two treatments. Viable regenerant plants were further cultivated on selective solid media.

Strains. Table 1 lists the nomenclature and derivation of all *P. patens* strains described in this paper.

Plasmids. Plasmid pLVneo103 was obtained from Prof. J. Schell (Hain et al. 1985). Plasmid pLVneo2103 (Hain et al. 1985), pABD1 (Paszowski et al. 1984), pHP23b (Paszowski et al. 1988) and pGL2 which contains a *Bam*HI fragment of the coding sequence of the gene *APH IV* (Gratz and Davies 1983), cloned in the polylinker of pDH51 (Pietrzak et al. 1986), were all kindly provided by Dr. J. Paszowski (ETH Zurich, Switzerland). pBR322 was obtained from Sigma Chemical Co. Plasmid DNA was isolated and purified using

Table 1. List of strains

Strain	Derivation or reference
Grandden wild-type	Ashton and Cove (1977)
Leiman wild-type	Isolated from the banks of Lac Lemun, Switzerland, 1982
pabA3	Ashton and Cove (1977)
nicA4	Ashton and Cove (1977)
nk85/y66	Ashton and Cove (1977)
c-7	pabA3 transformed by pLVneo103, selected for G418 resistance at 30 µg/ml
LS03	pabA3 transformed by pGL2, selected for hygromycin-resistance at 25 µg/ml
PHP 23 L2	Luman wild-type transformed by pHP23b, selected for G418 resistance at 50 µg/ml

standard procedures (Sambrook et al. 1989). Where applicable, DNA was linearised by restriction endonuclease digestion followed by phenol/chloroform extraction, ethanol precipitation and resuspension in 10 mM TRIS-HCl (pH 7.5)/1 mM EDTA (TE) at 1.0 mg/ml. Restriction enzymes were purchased from Bethesda Research Laboratories and used according to the suppliers' instructions. Calf thymus DNA was sheared to about 5–10 kb and used as carrier DNA.

Transformation procedure. The following modifications were made to the method described by Saul et al. (1988). Protoplasts were isolated and washed twice by centrifuging at 800 rpm for 5 min in 0.48 M mannitol. Protoplasts were counted and resuspended at 1.3×10^6 /ml in 0.48 M mannitol, 15 mM MgCl₂, 0.1% MES-KOH, pH 5.6 (MMM solution). Although the inclusion of W5 solution yielded transformants, it resulted in reduced viability and was routinely omitted. Protoplasts were heat shocked at 45°C for 5 min, cooled to 20°C and 300 µl (4×10^5 protoplasts) transferred into 10–14 ml sterile tubes. Thirty microlitres of DNA solution was added to each tube and gently mixed, followed by 300 µl of a solution of 40% (w/v) PEG 4000 (Prolabo) in 0.48 M mannitol containing 0.1 M Ca(NO₃)₂, pH 8.0. The PEG (polyethylene glycol) was either autoclaved before dissolving in the sterile mannitol solution or the final solution was sterilised by filtration. The PEG solution was left for 2–3 h before use and made fresh before each experiment.

Thirty microlitres of DNA solution routinely contained 5 µg of plasmid and 25 µg calf thymus carrier DNA (in Lausanne, 3 and 14 µg respectively) although 20–30 µg plasmid DNA only was also effective. The transformation mix was incubated at 20°C for 30 min with occasional gentle mixing. The PEG was diluted from the suspension by progressive step dilution over ca. 30 min with 10 ml of MMM solution. Protoplasts were centrifuged and resuspended in 0.3 ml MMM solution and, as previously described, either placed out in non-selective medium at a density of approximately 10^5 protoplasts per 9 cm petri dish or cultured in 6 ml liquid medium per 6 cm petri dish. Petri dishes were incubated in light (15 W/m²) for 4–7 days before transfer of the cellophane overlays to medium without mannitol and containing G418 (50 µg/ml) or hygromycin B (25–30 µg/ml). Antibiotic-resistant plants were observed and counted from 10 to 50 days after selection.

Plant DNA analysis. Genomic DNA was isolated from 1–5 g tissue (fresh weight, following drying on a filter by gentle vacuum suction) according to the miniprep procedure described by Dellaporta et al. (1983). This procedure yielded up to 50 µg DNA per gram of plant material. Transformant protonemata were grown on selective medium for at least 7 days, then washed twice in sterile water and filter-dried before freezing in liquid nitrogen. After digestion with restriction enzymes, the DNA was separated by electrophoresis in a 0.7% (w/v) agarose gel, denatured and transferred to nitrocellulose or nylon (Gene Screen Plus or Biodyne) membranes ac-

cording to standard procedures (Sambrook et al. 1989, or manufacturers' instructions) and hybridised with probes labelled with 32 P-dNTP by random hexamer priming (Feinberg and Vogelstein 1983). The genome size of *P. putans* has been estimated to be approximately 7×10^8 bp (N.H. Grimley and J.P. Fryd, unpublished data). In general, between 10^6 and 10^7 genomes were loaded per lane with an equivalent number of copies of probe DNA (see legends to the figures).

Crosses. Crosses were made according to Ashton et al. (1988) between an antibiotic-resistant *pubA3* transformant and the complementary auxotrophic and antibiotic-sensitive strain, *nicA4*. Where transformants had been isolated following treatment of the wild-type strain, crosses were made using the *nicB5/ylb6* strain, in which the *ylb* allele gives a yellow phenotype, to identify products of a cross rather than a self-fertilisation. The strains were co-inoculated in a sterile test tube and grown at 25°C for 3 weeks before transferring to 15°C for 2 weeks and then irrigating with sterile H_2O . In crosses between two auxotrophs, only spore capsules arising as a result of a cross should occur following irrigation with H_2O (Courtice et al. 1978).

Capsules, each containing about 5×10^5 viable spores, appeared after 3–4 weeks and were picked off and stored dry in a sterile Eppendorf tube at 4°C . Individual spore capsules were crushed in sterile H_2O and diluted aliquots

spread onto medium supplemented with both *p*-amino benzoic acid and nicotinic acid but without antibiotic. Protonemata from 100 sporelings were inoculated onto complete medium and grown for 10 days before inoculating fragments of each plant onto selective medium supplemented with *p*-amino benzoic acid and nicotinic acid, and non selective medium containing either one or the other vitamin. Plants were grown for 3 weeks before scoring for auxotrophies although segregation of antibiotic resistance was visible after 3 days. Resistant and sensitive progeny were selected and grown for DNA isolation and for Southern hybridisation analysis.

Results

Selection conditions

Protoplasts and protonemata of both wild types and *pubA3* were tested for their ability to grow on increasing concentrations of kanamycin sulphate, G418 and hygromycin B. On no occasions were surviving colonies detected after 10 days on G418 ($10 \mu\text{g/ml}$) and hygromycin B ($5 \mu\text{g/ml}$), whereas reduced growth still occurred on $100 \mu\text{g/ml}$ of kanamycin sulphate. We therefore decided to select for aminoglycoside phosphotransferase activity using the synthetic aminoglycoside G418 at $50 \mu\text{g/ml}$ and for hygromycin phosphotransferase activity using

hygromycin B at $25\text{--}30 \mu\text{g/ml}$. An additional advantage to using G418 rather than kanamycin was that G418 has a long life in illuminated culture, thus reducing the need for repeated subculture to maintain selection. Hygromycin degraded over time and cultures were usually subcultured within 2 to 3 weeks to maintain selection.

Primary selection

The transformation treatment reduces viability to about 10%. The frequencies of antibiotic-resistant plants growing at 10, 27 and 50 days after transfer to selection are presented in Table 2. The total number of stable transformants obtained from both experiments is also shown in Table 2. Controls included treatments where either no DNA or pBR322 DNA (vector-only control) was added and a treatment where the appropriate plasmid was added but no PEG was added. On no occasions were regenerants detected from these controls using selection levels of $50 \mu\text{g/ml}$ G418 or $25\text{--}30 \mu\text{g/ml}$ hygromycin B.

It can be seen from Table 2 that the initial transformation frequencies range between 1 and 84 regenerants per 10^4 viable protoplasts or 1 and 118 regenerants per μg DNA. However, the fate of these plants varies. Some regenerants grow strongly on selection and maintain resistance after a period of growth on non-selective media. These represent stable transformants and the frequency of occurrence of this class is low and variable, at best being equal to 10^{-4} viable protoplasts or 1 per μg DNA. Many regenerants do not grow beyond the 100 cells stage (usually reached by 10 days after transfer to selection) and it is likely that these cells have received multiple copies of the resistance gene which are only transiently expressed. A third class of regenerant continues to grow and may be subcultured and maintained for at least 2 years providing that selection is continuous. This class has been obtained frequently in both Leeds and Lausanne and we believe represents unstable transformation which may or may not involve integration (C.D. Knight and D. Schaefer, unpublished data).

It is possible that the structure of the plasmid also influences the transformation frequency and the class of clone recovered. Table 2 shows the number of regenerants obtained for some plasmids either in supercoiled or linear form. In all cases, supercoiled DNA yields a higher initial frequency of antibiotic-resistant regenerants; however, most of these are transient clones.

Mitotic and meiotic stability of stable transformants

The criteria of unrestricted growth on selection and maintenance of resistance after non-selective growth were used as indicators of stable transformation and mitotic stability. Protoplasts were isolated from protonemata of transformants HP 23 L2 and 15.03 which had been grown on non-selective and selective media. The protoplasts were regenerated on media with and without antibiotic and the regeneration frequencies found to be similar, regardless of the initial conditions of growth




Fig. 1. a–f. Each 5 cm plate was inoculated with 5 protonemata of either transformant c-7 (e–f) or the untransformed control *pubA3* (a, b) and grown for 14 days. At the start of the growth period all plates resembled plate b. Plates a and c do not contain antibiotic. Plates b, d, e and f contain G418 at $50 \mu\text{g/ml}$ (a, d and e) and $150 \mu\text{g/ml}$ (f). Plate e was inoculated with a culture grown for 14 days in the absence of selection.

(data not shown). A further indication of mitotic stability was noted by the successful fusion of protoplasts and selection of fertile diploid doubly-resistant clones (D. Schaefer, unpublished data). The characteristics of one stable transformant (c-7) are shown in Fig. 1. Transformant c-7 was selected on medium containing G418 at $50 \mu\text{g/ml}$ but was able to grow at $150 \mu\text{g/ml}$ G418, albeit with some restriction when compared with the untransformed control under non-selective conditions. The same transformant was tested for its ability to grow on $50 \mu\text{g/ml}$ after a period of 14 days growth in the absence of selection. Figure 1 shows that this treatment does not affect the growth pattern of transformant c-7.

Hygromycin- and G418-resistant clones were taken through meiosis on selective and non-selective media and spores were tested for their ability to germinate on medium containing antibiotics. A number of different strains stably transformed for either G418 or hygromycin resistance have been shown to transmit resistance to 100% of spores through up to four generations (data not shown).

DNA analysis

Southern hybridisation analysis of a stable transformant revealed the presence of multiple copies of the plasmid DNA integrated at a single site in the nuclear genome. Figure 2 shows the hybridisation pattern of DNA from the G418-resistant transformant c-7, probed with the 2.2 kb *EcoRI*-*Sall* fragment of the transforming plasmid pLGVneo103 which contains the *NPT-II* gene (Hain et al. 1985). The quantities of DNA loaded per lane are

Table 2. Relative transformation frequencies

Plasmid (promoter)	Plasmid ^a Structure	Selection	Frequency of plants growing (days after transfer to selection)	Total number of stable transformants
pLGVneo103 (neo)	§	G418	17 13 37 30 NC NC	1
pLGVneo203 (neo)	§	G418	12 17 NC NC	9 13 0
pABD1 ⁴ (CaMV 19S)	§	G418	75 190 24 32 2 2	0
pABD1 ⁴ (CaMV 19S)	¶	G418	1 1 1 1 1 1	0
pIP23b (CaMV 35S)	§	G418	84 118 80 112 16 23 2	2
pIP23b (CaMV 35S)	¶	G418	23 25 42 46 24 27 2	2
pGL2 ⁴ (CaMV 35S)	§	hyg	35 49 13 18 13 18 0	0
pGL2 (CaMV 35S)	¶	hyg	6 8 12 16 12 15 3	3
No plasmid		G418:hyg	0 0 0 0 0 0 0	0
pBR322 (vector only)	§ and ¶	G418:hyg	0 0 0 0 0 0 0	0
No PEG	§ and ¶	G418:hyg	0 0 0 0 0 0 0	0

NC, not counted

a. Frequency per 10^4 viable protoplasts; b. frequency per μg DNA

§. Supercoiled (s); ¶. linear (l)

⁴ Data from one experiment only, all other data are the mean of two experiments

Analysis of conserved domains identifies a unique structural feature of a chloroplast heat shock protein

Qiang Chen and Elizabeth Vierling

Department of Biochemistry, University of Arizona, Tucson, AZ 85721, USA

Received August 6, 1990

Introduction

Summary. A low molecular weight heat shock protein which localizes to chloroplasts has been identified in several plant species. This protein belongs to a eukaryotic superfamily of small HSPs, all of which contain a conserved carboxyl-terminal domain. To investigate further the structure of this HSP, we isolated and sequenced cDNA clones for the chloroplast LMW HSPs from *Petunia hybrida* and *Arabidopsis thaliana*. The cloning of chloroplast HSPs from these two species enabled us to compare the amino acid sequences of this protein from plant species (*Petunia*, *Arabidopsis*, pea, soybean and maize) that represent evolutionarily divergent taxonomic subclasses. Three conserved regions were identified, which are designated as regions I, II and III. Regions I and II are also shared by cytoplasmic LMW HSPs and therefore are likely to have functional roles common to all eukaryotic LMW HSPs. In contrast, consensus region III is not found in other LMW HSPs. Secondary structure analysis predicts that this region forms an amphipathic α -helix with high conservation of methionine residues on the hydrophobic face and 100% conservation of residues on the hydrophilic face. This structure is similar to three helices, termed "methionine bristles", which are found in a methionine-rich domain of a 54 kDa protein component of signal recognition particle (SRP54). The conservation of regions I and II among LMW cytoplasmic and chloroplast HSPs suggests that these HSPs perform related functions in different cellular compartments. However, identification of the methionine bristle domain suggests that chloroplast HSPs also have unique functions or substrates within the special environment of the chloroplast or other plastids.

Key words: Small HSPs – Protein transport – *Arabidopsis* – Methionine bristle – Amphipathic helix

Abbreviations: HS, heat shock; HSP, heat shock protein; LMW, low molecular weight

Offprint requests to: E. Vierling

- Jenkins GI, Cove DJ (1983b) Phototropism and polarotropism of primary chloronema of the moss, *Physcomitrella patens*: responses of the wild-type. *Planta* 158:337–364
- Knight CD, Cove DJ, Boyd PJ, Ashton NW (1988) The isolation of biochemical and developmental mutants in *Physcomitrella patens*. In: Cline JM (ed) Methods in bryology. Proceedings of Bryology Methods Workshop, Mainz Botanical Laboratory, Nidhagen, pp 47–58
- Kochler F, Cardon G, Pochlman M, Gill R, Schieder O (1989) Enhancement of transformation rates in higher plants by low dose irradiation: Are DNA repair systems involved in the incorporation of exogenous DNA into the plant genomes? *Plant Mol Biol* 12:165–199
- Long Z, Wang S-Y, Nelson N (1989) Cloning and nucleotide sequence analysis of gene coding for the major chlorophyll-binding protein of the moss *Physcomitrella patens* and the halobacterium *Halobacterium salinarum*. *Gene* 76:295–312
- Meyer P, Karske S, Niedebohn I, Heldmann I, Busman K, Siedler H (1989) A genomic DNA segment from *Petunia hybrida* leads to increased transformation frequencies and simple integration patterns. *Proc Natl Acad Sci USA* 85:3568–3572
- Meyrowitz EM (1989) *Arabidopsis*, a useful weed. *Cell* 56:263–269
- Pazkowski J, Shillito RD, Saul M, Mandak V, Hohn T, Hohn B, Potrykus I (1984) Direct gene transfer to plants. *EMBO J* 3:2717–2723
- Pazkowski J, Baur M, Bogucki A, Potrykus I (1988) Gene targeting in plants. *EMBO J* 7:4021–4026
- Pieratz M, Shillito RD, Hohn T, Potrykus I (1986) Expression in plants of two bacterial antibiotic resistance genes after protoplast transformation with a new plant expression vector. *Nucleic Acids Res* 14:3857–3868
- Sambrook J, Fritsch EF, Maniatis T (1989) Molecular cloning. A laboratory manual, 2nd edition. Cold Spring Harbour Laboratory, Cold Spring Harbor, NY
- Saul MW, Shillito RD, Negrutiu I (1988) Direct DNA transfer to protoplasts with and without electroporation. In: Gelvin SB, Schilperhoort RA (eds) Plant molecular biology manual. Kluwer Academic Publishers, Netherlands, pp 1–16
- Scherdin U, Rhodes K, Breidl M (1990) Transcriptionally active genome regions are preferred targets for retrovirus integration. *J Virol* 64:907–912
- Shillito R, Pazkowski J, Potrykus I (1983) Agarose plating and bead-type culture technique enable and stimulate development of protoplast-derived colonies in a number of plant species. *Plant Cell Report* 2:244–247
- Snell WJ (1983) Cell-cell interactions in *Chlamydomonas*. *Annu Rev Plant Physiol* 36:287–315

Communicated by H. Siedler

- Quile PJ, Hughes KW (1987) The consensus land plant chloroplast gene order is present, with two alterations, in the moss *Physcomitrella patens*. *Mol Gen Genet* 208:335–341
- Courtice GRM, Ashton NW, Cove DJ (1978) Evidence for the restricted passage of metabolites into the sporophyte of the moss *Physcomitrella patens*. *J Biol* 10:191–198
- Cove DJ, Ashton NW (1984) The hormonal regulation of gametophyte development in bryophytes. In: Dyer AF, Duckett JG (eds) Ecophysiology of bryophytes. Academic Press, London, pp 177–201
- Cove DJ, Schild A, Ashton NW, Hartmann E (1978) Genetic and physiological studies of the effect of light on the development of the moss *Physcomitrella patens*. *Photochem Photobiol* 27:249–254
- Cove DJ, Kammerer W, Knight CD, Leeth MJ, Martin CR, Wang TL (1990) Developmental genetic studies of the moss, *Physcomitrella patens*. In: Jenkins GI, Schuch W (eds) Molecular biology of plant development. Company of Biologists, Cambridge, UK (in press)
- Crispen D, Shull S (1982) Regulation of DNA ligase activity by poly(ADP-ribose). *Nature* 296:271–272
- DellaPorta SL, Wood J, Hicks JB (1983) A plant DNA miniprepation. *Version II*. *Plant Mol Biol Report* 1:19–21
- Faithorne DR, Cove DJ, Ashton NW (1990) Genetic analysis by somatic hybridization of cytokinin overproducing developmental mutants of the moss *Physcomitrella patens*. *Mol Gen Genet* 222:217–224
- Faithorne AP, Vegetarian B (1983) A technique for radiolabelling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* 132:6–13
- Giardi-Chiari GB, Cherdronswart W, Dewulf J, Pazkowski J, Jacobs M, Negrutiu I (1990) Hybrid genes in the analysis of transformation conditions. 3. Temporal/spatial fate of *NPTII* gene integration, its inheritance and factors affecting these processes in *Nicotiana glauca*. *Plant Mol Biol* 14:687–696
- Grimsey NH, Ashton NW, Cove DJ (1977a) The production of somatic hybrids by protoplast fusion in the moss *Physcomitrella patens*. *Mol Gen Genet* 154:97–100
- Grimsey NH, Ashton NW, Cove DJ (1977b) Complementation analysis of autotrophic mutants of the moss *Physcomitrella patens*, using protoplast fusion. *Mol Gen Genet* 155:103–107
- Gritz L, Davies J (1983) Plasmid encoded hygromycin resistance: the sequence of hygromycin B phosphotransferase gene and its expression in *Escherichia coli* and *Saccharomyces cerevisiae*. *Gene* 25:179–188
- Hahn R, Subel P, Camillio AP, Sieschins HH, Herren-Eustalla L, Schell J (1983) Uptake, integration, expression and genetic transmission of a selectable chimeric gene by plant protoplasts. *Mol Gen Genet* 197:161–168
- Jenkins GI, Cove DJ (1983a) Light requirements for regeneration of the moss, *Physcomitrella patens*. *Planta* 157:39–45

Note added in proof

The transformation procedure has been improved since the manuscript was submitted. A modified protocol is available upon request.